

Exhibit A

U.S. Patent 8,246,979 B2



US008246979B2

(12) **United States Patent**
Schacht et al.

(10) **Patent No.:** **US 8,246,979 B2**
(45) **Date of Patent:** **Aug. 21, 2012**

- (54) **TRANSDERMAL DELIVERY SYSTEM FOR THE ADMINISTRATION OF ROTIGOTINE**
- (75) Inventors: **Dietrich Wilhelm Schacht**, Köln (DE); **Mike Hannay**, Wachtberg-Villprott (DE); **Hans-Michael Wolff**, Monheim (DE)
- (73) Assignee: **UCB Pharma GmbH**, Monheim (DE)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 1502 days.
- (21) Appl. No.: **10/623,864**
- (22) Filed: **Jul. 22, 2003**
- (65) **Prior Publication Data**
US 2005/0079206 A1 Apr. 14, 2005
- Related U.S. Application Data**
- (60) Provisional application No. 60/451,716, filed on Mar. 5, 2003.
- (30) **Foreign Application Priority Data**
Jul. 30, 2002 (EP) 02016860

- (51) **Int. Cl.**
A61F 13/00 (2006.01)
A61F 13/02 (2006.01)
A61K 9/70 (2006.01)
A61L 15/16 (2006.01)
- (52) **U.S. Cl.** 424/449; 424/443; 424/447; 424/448
- (58) **Field of Classification Search** 424/449, 424/443, 447, 448
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,797,494 A	3/1974	Zaffaroni	424/434
4,588,580 A	5/1986	Gale et al.	424/449
4,915,950 A	4/1990	Miranda et al.	424/448
4,973,468 A	11/1990	Chiang et al.	424/449
5,069,909 A	12/1991	Sharma et al.	424/449
5,091,186 A	2/1992	Miranda et al.	424/448
5,124,157 A	6/1992	Colley et al.	424/448
5,147,916 A	9/1992	Sweet	524/266
5,177,112 A	1/1993	Horn	514/65
5,217,718 A	6/1993	Colley et al.	424/449
5,225,198 A	7/1993	Sharma et al.	424/443
5,232,702 A *	8/1993	Pfister et al.	424/448
5,234,690 A	8/1993	Chiang et al.	424/448
5,246,997 A	9/1993	Sweet	524/266
5,252,334 A	10/1993	Chiang et al.	424/448
5,252,335 A	10/1993	Chiang	424/449
5,271,940 A	12/1993	Cleary et al.	424/448
5,273,755 A	12/1993	Venktraman et al.	424/448
5,273,756 A	12/1993	Fallon et al.	424/448
5,273,757 A	12/1993	Jaeger et al.	424/448
5,308,625 A	5/1994	Wong et al.	424/449
5,382,596 A	1/1995	Sleeve et al.	514/459
5,393,529 A	2/1995	Hoffmann et al.	424/445
5,456,745 A	10/1995	Roreger et al.	106/128
5,500,222 A	3/1996	Lee et al.	424/448

5,554,381 A	9/1996	Roos et al.	424/449
5,601,839 A	2/1997	Quan et al.	424/448
5,670,164 A	9/1997	Meconi et al.	424/448
5,676,968 A	10/1997	Lipp et al.	424/448
5,688,524 A	11/1997	Hsu et al.	424/449
5,733,571 A *	3/1998	Sackler	424/449
5,771,890 A	6/1998	Tamada	128/635
5,788,983 A *	8/1998	Chien et al.	424/449
5,807,570 A	9/1998	Chen et al.	424/449
5,834,010 A	11/1998	Quan et al.	424/448
5,840,336 A	11/1998	Hsu et al.	424/484
5,876,746 A	3/1999	Jona et al.	424/449
5,879,701 A	3/1999	Audett et al.	424/448
5,891,461 A	4/1999	Jona et al.	424/449
5,902,603 A	5/1999	Chen et al.	424/449
5,906,830 A	5/1999	Farinas et al.	424/448
5,932,240 A	8/1999	D'Angelo et al.	424/449
5,980,932 A	11/1999	Chiang et al.	
6,024,974 A	2/2000	Li	424/448
6,024,976 A	2/2000	Miranda et al.	424/449
6,063,398 A	5/2000	Gueret	424/443
6,218,421 B1	4/2001	King	514/421
6,299,900 B1	10/2001	Reed et al.	
6,316,022 B1 *	11/2001	Mantelle et al.	424/448
6,372,920 B1	4/2002	Minaskanian et al.	549/75
6,393,318 B1	5/2002	Conn et al.	604/20
6,398,562 B1	6/2002	Butler et al.	439/91
6,465,004 B1 *	10/2002	Rossi-Montero et al.	424/448
6,514,530 B2 *	2/2003	Skluzacek et al.	424/468
6,620,429 B1	9/2003	Müller	

(Continued)

FOREIGN PATENT DOCUMENTS

CA 2 532 804 2/2005

(Continued)

OTHER PUBLICATIONS

Blindauer (2003) Arch. Neurol. 60(12): 1721-1728.
Chiang et al. (1995) Proc. Int. Symp. Controlled Release Bioact. Mater. 22, 710-711.
H.F. Hammond in D. Satas Handbook of Pressure Sensitive Adhesive Technology (1989) 2nd Ed. Chapter 4, p. 38.
Hsu et al. (1992) Cygnus Therapeutic Systems Project Report N-0923, 2-19.
Levien et al. (2005) Advances in Pharmacy 3(1): 62-92.
LeWitt et al. (2007) Neurology 68, 1262-1267.
Löschnmann et al. (1989) Eur. J. Pharmacol. 166: 373-380.
Pfister (1988) Drug and Cosmetic Ind. (Oct): 44-52.
Pfister (1989) Pharm. Tech. (March): 126-138.
Pfister and Hsieh (1990) Pharm. Tech. (Sept): 132-140.
Pfister and Hsieh (1990) Pharm. Tech. (Oct): 54-60.
Pfister et al. (1991) Chemistry in Britain (Jan): 43-46.
Pfister et al. (1992) Pharm. Tech. (Jan): 42-58 and 83.
Roy et al. (1996) J. Pharm. Science 85(5): 491-495.

(Continued)

Primary Examiner — Richard Schnizer

Assistant Examiner — Audrea Buckley

(74) Attorney, Agent, or Firm — Harness, Dickey & Pierce, P.L.C.

(57) **ABSTRACT**

An improved transdermal delivery system (TDS) comprises a self-adhesive matrix comprising a solid or semi-solid semi-permeable polymer which contains rotigotine in its free base form as a multitude of microreservoirs within the matrix. The self-adhesive matrix is highly permeable to the free base of rotigotine and is impermeable to the protonated form of rotigotine.

18 Claims, 7 Drawing Sheets

US 8,246,979 B2

Page 2

U.S. PATENT DOCUMENTS

6,685,959	B1	2/2004	Moreau et al.	424/449
6,687,522	B2	2/2004	Tamada	600/347
6,699,498	B1	3/2004	Müller	424/449
6,884,434	B1 *	4/2005	Muller et al.	424/487
6,899,894	B1 *	5/2005	Klein et al.	424/448
6,929,901	B2 *	8/2005	Lui et al.	430/329
7,279,178	B1	10/2007	Berthold	424/443
7,309,497	B2	12/2007	Rimpler et al.	424/422
7,413,747	B2	8/2008	Mueller et al.	424/448
2001/0053777	A1 *	12/2001	Brecht	514/221
2002/0004065	A1	1/2002	Kanios	424/449
2002/0110585	A1	8/2002	Godbey et al.	424/449
2003/0026830	A1	2/2003	Lauterback et al.	424/449
2003/0027793	A1	2/2003	Lauterback et al.	514/63
2003/0065086	A1 *	4/2003	Kosal	524/588
2003/0166709	A1	9/2003	Rimpler et al.	514/447
2004/0013620	A1	1/2004	Klose et al.	424/59
2004/0048779	A1	3/2004	Schollmayer	514/2
2004/0057985	A1	3/2004	Bracht	424/449
2004/0081683	A1	4/2004	Schacht et al.	424/449
2004/0110673	A1	6/2004	Steinkasserer et al.	514/12
2004/0116537	A1	6/2004	Li et al.	514/663
2004/0137045	A1	7/2004	Breitenbach et al.	424/449
2005/0033065	A1	2/2005	Mueller et al.	549/74
2005/0079206	A1	4/2005	Schacht et al.	424/449
2005/0136101	A1	6/2005	Berthold	424/448
2005/0175678	A1	8/2005	Breitenbach et al.	424/448
2005/0175680	A1	8/2005	Morgan et al.	424/449
2005/0197385	A1	9/2005	Scheller et al.	514/438
2005/0260254	A1	11/2005	Breitenbach et al.	424/449
2006/0222691	A1	10/2006	Cantor et al.	424/448
2006/0263419	A1	11/2006	Wolff	424/448
2007/0072917	A1	3/2007	Scheller et al.	514/357
2007/0093546	A1	4/2007	Scheller et al.	514/447
2007/0191308	A1	8/2007	Kramer	514/60
2007/0191470	A1	8/2007	Scheller	514/438
2007/0197480	A1	8/2007	Scheller et al.	514/114
2008/0008748	A1	1/2008	Beyreuther et al.	424/449
2008/0138389	A1	6/2008	Muller et al.	424/448
2008/0146622	A1	6/2008	Scheller et al.	514/357
2008/0274061	A1	11/2008	Schollmayer et al.	424/45
2009/0143460	A1	6/2009	Wolff et al.	514/438
2010/0311806	A1	12/2010	Wolff et al.	514/438

FOREIGN PATENT DOCUMENTS

CA	2 532 859	2/2005
CA	2 547 820	6/2005
CA	2 546 797	7/2005
DE	198 14 083	10/1999
DE	199 40 238	3/2001
DE	10060550	4/2002
EP	1 256 340	11/2002
FR	2792529	10/2000
WO	WO 93/07842	4/1993
WO	WO 93/14727	8/1993
WO	WO 93/16073	8/1993
WO	WO 94/04109	3/1994
WO	WO 94/07468	4/1994
WO	WO 95/00122	1/1995
WO	WO 95/01767	1/1995
WO	WO 95/05137	2/1995
WO	WO 95/05138	2/1995
WO	WO 96/00110	1/1996
WO	WO 96/22083	7/1996
WO	WO 96/22084	7/1996
WO	WO 96/40087	12/1996
WO	WO 97/09971	3/1997
WO	WO 97/29735	8/1997
WO	WO 98/00126	1/1998
WO	WO 99/01795	1/1999
WO	WO 99/49852	10/1999
WO	WO 02/26217	4/2002
WO	WO 02/045699	6/2002
WO	WO 02/074286	9/2002
WO	WO 02/089777	11/2002

WO	WO 2004/012719	2/2004
WO	WO 2004/012730	2/2004
WO	WO 2004/058247	6/2004

OTHER PUBLICATIONS

Tanojo et al. (1997) J. controlled Release 45, 41-47.
 Thomas et al. (1991) STP Pharma Sci 1(1): 38-46.
 Tozer (1995) Clinical Pharmacokinetics, 480-483.
 Ulman et al. (1998) Information Brochure, The Correlation of Tape Properties and Rheology, Dow Corning Group.
 The United States Pharmacopeia/The National Formulary (2002) USP25/NF20, Chapter 724.
 Office Action, dated Feb. 2, 2009 issued in U.S. Appl. No. 10/627,990.
 Office Action, dated Aug. 18, 2009 issued in U.S. Appl. No. 10/627,990.
 Office Action, dated Apr. 27, 2010 issued in U.S. Appl. No. 10/627,990.
 Office Action, dated Mar. 23, 2007 issued in U.S. Appl. No. 10/713,424.
 Office Action, dated Sep. 14, 2007 issued in U.S. Appl. No. 10/713,424.
 Office Action, dated Jun. 25, 2008 issued in U.S. Appl. No. 10/713,424.
 Office Action, dated Feb. 19, 2009 issued in U.S. Appl. No. 10/713,424.
 Office Action, dated Sep. 13, 2007 issued in U.S. Appl. No. 10/936,620.
 Office Action, dated May 1, 2008 issued in U.S. Appl. No. 10/936,620.
 Office Action, dated Jan. 26, 2009 issued in U.S. Appl. No. 10/936,620.
 Office Action, dated Nov. 6, 2009 issued in U.S. Appl. No. 10/936,620.
 Office Action, dated Sep. 21, 2006 issued in U.S. Appl. No. 10/429,283.
 Office Action, dated Jan. 8, 2008 issued in U.S. Appl. No. 10/429,283.
 Office Action, dated Sep. 12, 2008 issued in U.S. Appl. No. 10/429,283.
 Office Action, dated Mar. 30, 2009 issued in U.S. Appl. No. 10/429,283.
 Office Action, dated Dec. 23, 2009 issued in U.S. Appl. No. 10/429,283.
 Office Action, dated Apr. 15, 2002 issued in U.S. Appl. No. 09/647,290.
 Office Action, dated Nov. 1, 2002 issued in U.S. Appl. No. 09/647,290.
 Office Action, dated Aug. 6, 2008 issued in U.S. Appl. No. 10/517,157.
 Office Action, dated Mar. 6, 2009 issued in U.S. Appl. No. 10/517,157.
 Office Action, dated Nov. 24, 2009 issued in U.S. Appl. No. 10/517,157.
 Office Action, dated Jun. 9, 2010 issued in U.S. Appl. No. 11/931,762.
 Office Action, dated Jan. 23, 2008 issued in U.S. Appl. No. 11/931,666.
 Office Action, dated Oct. 15, 2009 issued in U.S. Appl. No. 10/139,894.
 Goodman and Gilman's, (1990) The Pharmacological Basis of Therapeutics, Pergamon Press, 8th ed, p. 160.
 The Merck Index, Thirteenth Edition (2001) "7024. Oxybutynin"
 Whitehouse Station, NJ: Merck & Co., Inc., p. 1245.
 Miyamoto et al. (1994) Analyst 119, 1489-1492.
 Nugroho et al. (2004) Pharmaceutical Research, 844-850.
 Roy (1990) "Transdermal delivery of narcotic analgesics: PH, anatomical, and subject influences on cutaneous permeability of fentanyl and sufentanil." Pharmaceutical Research, New York, NY, US, vol. 7, No. 8, pp. 842-847, XP002923066.
 Stryer, (1996) Biochemistry, W.H. Freeman and Company, New York, pp. 20-22 and 25.

US 8,246,979 B2

Page 3

<http://www.accessdata.fda.gov> (USFDA label for NEUPRO (2007) retrieved by CA examiner from Drugs@FDA database).

<http://www.neupro.com> (Neupro recall letter to physicians (2008), retrieved from CA examiner from US NEUPRO homepage, attached).

Office Action, dated Oct. 8, 2010 issued in U.S. Appl. No. 10/936,620.

Office Action, dated Jan. 4, 2011 issued in U.S. Appl. No. 10/627,990.

Office Action, dated Aug. 3, 2010 issued in U.S. Appl. No. 10/517,157.

Office Action, dated Nov. 9, 2010 issued in U.S. Appl. No. 12/114,348.

Office Action, dated Oct. 1, 2010 issued in U.S. Appl. No. 10/429,283.

* cited by examiner

U.S. Patent

Aug. 21, 2012

Sheet 1 of 7

US 8,246,979 B2

Fig. 1




Fig. 2





Fig. 3




U.S. Patent

Aug. 21, 2012

Sheet 4 of 7

US 8,246,979 B2

Fig. 4



U.S. Patent

Aug. 21, 2012

Sheet 5 of 7

US 8,246,979 B2

Fig. 5




Fig. 6





Fig. 7



US 8,246,979 B2

1

**TRANSDERMAL DELIVERY SYSTEM FOR
THE ADMINISTRATION OF ROTIGOTINE**

This application claims benefit of U.S. Provisional Patent Application No. 60/451,716 filed Mar. 5, 2003, the entire contents of which are hereby incorporated by reference in their entirety.

This application contains subject matter that is related to U.S. patent application Ser. No. 10/627,990, filed on Jul. 28, 2003.

FIELD OF INVENTION

The present invention relates to an improved transdermal delivery system for rotigotine. Moreover, the invention relates to a method of treatment using the transdermal delivery system.

TECHNICAL BACKGROUND

To date, various transdermal delivery systems (TDS) for the administration of rotigotine have been described. WO 94/07468 discloses a TDS containing rotigotine hydrochloride as active substance in a two-phase matrix, which is essentially formed by a hydrophobic polymer material as the continuous phase and a disperse hydrophilic phase contained therein and mainly containing the drug and hydrated silica. The silica is said to enhance the maximum possible loading of the TDS with the hydrophilic salt. Moreover, the formulation of WO 94/07468 usually contains additional hydrophobic solvents, permeation promoting substances, dispersing agents and, in particular, an emulsifier which is required to emulsify the aqueous solution of the active component in the lipophilic polymer phase. A TDS prepared by using such a system has been tested in healthy subjects and Parkinson's patients. However, no satisfactory drug plasma levels were achieved.

Various further TDSs have been described in WO 99/49852.

In the TDS according to WO 94/07468 and many related applications, passive diffusion membranes were used.

However, as the skin is to be seen as a very efficient barrier for most drug candidates, such type of membrane controlled systems are more or less limited in practice to transdermal delivery of active substances that reveal a very high skin permeability. Additionally, special requirements on drug release kinetics have to be met like contact delivery over several days.

An object of the present invention is to control (i.e. to canalise/manoeuvre) the transport of rotigotine towards and across the skin from a drug reservoir, thereby enhancing the flux of rotigotine across the TDS/skin interface.

A further object and aspect of the present invention is to provide a suitable composition and manufacturing methods of polymer matrices in TDS which lead to an enhanced delivery of rotigotine to and across the skin by

- (i) preventing back diffusion of the drug portion which is ionized in the skin according to its pKa value—from the skin tissue into the TDS,
- (ii) offering continuous delivery of the active compound across the stratum corneum not only via the common more lipophilic route (e.g. intercellular) but also through hydrophilic pores (e.g. eccrine sweat glands).

SUMMARY OF THE INVENTION

There is now provided a TDS comprising a backing layer inert to the components of the matrix, a self-adhesive matrix containing rotigotine, and a protective foil or sheet to be removed prior to use,

2

characterized in that the self-adhesive matrix comprises a solid or semi-solid semi-adhesive polymer

- (1) wherein rotigotine in its free base form has been incorporated,
- (2) which comprises a multitude of microreservoirs within the matrix, said microreservoirs containing rotigotine and optionally at least a crystallization inhibitor,
- (3) which is permeable to the free base of rotigotine,
- (4) which is substantially impermeable to the protonated form of rotigotine,
- (5) wherein the maximum diameter of the microreservoirs is less than the thickness of the matrix.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows the effect of the protonation of rotigotine in the semi-permeable matrix on the drug absorption.

FIG. 2 shows the impact of the size distribution of the microreservoirs in the semi-permeable matrix on the drug absorption.

FIG. 3 shows the effect of reducing the amount of the protonated form of rotigotine in the semi-permeable matrix and reducing the size of the microreservoirs on the drug absorption.

FIG. 4 shows a microscope image of a conventional TDS.

FIG. 5 shows a microscope image of the TDS according to the invention.

FIG. 6 shows the effect of reducing the amount of the protonated form of rotigotine in the semi-permeable matrix and reducing the size of the microreservoirs on the in vitro skin permeation of the drug.

FIG. 7 shows a comparison of the in vitro skin permeation of rotigotine for the TDS of the invention and an acrylate-based TDS.

DESCRIPTION OF THE INVENTION

The present invention provides a TDS providing a high steady state flux rate of rotigotine over the TDS/skin interface.

Surprisingly, it was found that drug release properties of a TDS having a silicone-type adhesive matrix containing rotigotine can be significantly enhanced by

- (1) minimizing the amount of rotigotine which is present in the protonated form (salt form);
- (2) incorporating rotigotine in a multitude of microreservoirs within the self-adhesive matrix consisting of a solid or semi-solid semi-permeable polymer.

The impact of above described measures on drug release characteristics of rotigotine in vivo is illustrated in FIGS. 1, 2 and 3. The relative drug absorption in vivo was highest for the sample according to the invention; increase of the size of the microreservoirs and/or the amount of drug salt residues in the TDS led to slower initial drug release. Based on the above findings, the present invention was accomplished.

When using the TDS, according to the present invention, a high transfer of rotigotine from the silicone matrix into the outermost skin layer can be achieved. Consequently, plasma values of rotigotine are sufficient to allow for a reasonable expectation that an efficient treatment with fewer side effects can be provided.

The drug contained in the TDS according to the invention is 5,6,7,8-tetrahydro-6-[propyl-[2-(2-thienyl)ethyl]amino]-1-naphthalenol (INN: rotigotine). Rotigotine is a dopamine D2 receptor antagonist, which is useful for example in the treatment of Parkinson's disease.

US 8,246,979 B2

3

It should be understood that the term "treatment" in the context of this application is meant to designate a treatment or an alleviation of the symptoms. The treatment may be of a therapeutic or prophylactic nature.

It will be understood by a person skilled in the art that rotigotine exists in various isomeric forms. It has to be understood that any single isomer or a mixture of different isomers may be used in the TDS according to the invention. Hence, the S- or R-enantiomer or the racemate or any other enantiomer mixture of rotigotine may be used.

At least a part of rotigotine is contained in a multitude of microreservoirs distributed within the self-adhesive matrix of the TDS according to the invention. This does not exclude and will normally even imply that a certain fraction of rotigotine is dissolved in the solid or semi-solid semi-permeable polymer of the matrix at its saturation concentration.

Within this specification "microreservoirs" are meant to be understood as particulate, spatially and functionally separate compartments consisting of pure drug or a mixture of drug and crystallization inhibitor, which are dispersed in the self-adhesive (polymer) matrix. Preferably the self-adhesive matrix contains 10^3 to 10^9 microreservoirs per cm^2 of its surface, particularly preferred are 10^6 to 10^9 microreservoirs per cm^2 .

Rotigotine is incorporated in the self-adhesive matrix in its free base form. This does not totally exclude the presence of some residual salt form of rotigotine in the final TDS. However, the salt form of rotigotine should be contained in the self-adhesive matrix of the final TDS in an amount of preferably less than 5%, more preferably less than 2%, particularly less than 1% (w/w).

If rotigotine is present in the self-adhesive matrix in its protonated (salt) form, it will not be released by the self-adhesive matrix. Thus, the amount of the salt form of rotigotine can be determined by performing a drug dissolution test according to the Paddle over Disk method as described in the United States Pharmacopeia (United States Pharmacopeia/New Formulary (USP25/NF20), Chapter 724 "Drug Release", United States Pharmacopeial Convention, Inc., Rockville, Md. 20852, USA (2002)), using the following conditions: dissolution medium: 900 ml phosphate buffer pH 4.5; temperature adjusted to $32 \pm 0.5^\circ \text{C}$; paddle rotation speed: 50 rpm; sampling times: 0.5, 1, 2 and 3 h, respectively. The increase in the eluted rotigotine concentration can be used to calculate the amount of unprotonated rotigotine in the matrix.

The amount of the salt form of rotigotine may be reduced e.g. by reducing the water content of the mass containing the drug and organic solvent(s). In a particularly preferred embodiment of the invention the water content is reduced during manufacture to preferably less than 0.4% (w/w), more preferably less than 0.1%, of the mass.

A further step, which may be taken for reducing the amount of the salt form of rotigotine, is isolating the free base form of rotigotine in solid form prior to the preparation of the TDS. If the free base of rotigotine is produced in situ during the manufacture of the TDS by neutralizing an acid addition salt, a certain residue of the ionized drug form will remain in the polymer matrix (usually $>5\%$ (w/w) and up to approximately 10%). Therefore, such in situ preparation of the free base form will generally not be suitable for practising the present invention.

The maximum diameter of the microreservoirs is less than the thickness of the matrix, preferably up to 70% of the thickness of the matrix, particularly preferably 5 to 60% of the thickness of the matrix. For an exemplary thickness of the matrix of 50 μm this corresponds to a maximum diameter of

4

the microreservoirs in the range of preferably up to 35 μm . The term "maximum diameter" is meant to be understood as the diameter of the microreservoirs in one dimension (x-, y-, or z-dimension), which is the largest. It is clear to the skilled person that in case of spherical diameters the maximum diameter corresponds to the microreservoir's diameter. However, in the case of microreservoirs, which are not shaped in the form of spheres—i.e. of different geometric forms—the x-, y- and z-dimensions may vary greatly.

As the maximum diameter of the microreservoirs in the direction of the cross-section of the matrix, i.e. between the release surface and the backing layer, is less than the thickness of the matrix, direct contact between the skin and the basic microreservoirs containing rotigotine is avoided, if not at all prevented. Owing to the slightly acidic pH of the skin, direct contact between the skin and the microreservoirs in the matrix leads to protonation of rotigotine, thereby deteriorating the semi-permeability of the matrix.

In a particularly preferred embodiment of the invention, the mean diameter of the microreservoirs containing rotigotine distributed in the matrix is in the range of 1 to 40%, even more preferably 1 to 20%, of the thickness of the drug-loaded self-adhesive matrix. For an exemplary thickness of the matrix of 50 μm this corresponds to a mean diameter of the microreservoirs in the range of preferably 0.5 to 20 μm . The term "mean diameter" is defined as the mean value of the x,y,z-average diameters of all microreservoirs. The target particle size can be adjusted by the solids content and viscosity of the drug-containing coating mass.

The maximum and mean diameters of the microreservoirs as well as the number of microreservoirs per surface area of the self-adhesive matrix can be determined as follows: The release liner is removed from the TDS, and the free adhesive surface is examined with a light microscope (Leica microscope type DM/RBE equipped with a camera type Basler A 113C). The measurement is performed by incidental polarized light analysis using a microscope at 200 \times magnification. A picture analysis is performed using the software Nikon Lucia_Di, Version 4.21, resulting in mean and maximum diameters for each sample.

The TDS of the present invention is of the "matrix" type. In such matrix type TDS the drug is dispersed in a polymer layer. The TDS of the matrix type in their simplest version comprise a one-phase (monolayer) matrix. They consist of a backing layer, a self-adhesive matrix containing the active agent and a protective foil or sheet, which is removed before use.

Versions that are more complicated comprise multi-layer matrixes, wherein the drug may be contained in one or more non-adhesive polymer layers. The TDS of the present invention is preferably a one-phase (mono layer) matrix system.

The solid or semi-solid semi-permeable polymer of the self-adhesive matrix has to satisfy the following requirements:

1. Sufficient solubility and permeability for the free base form of rotigotine.
2. Impermeability for the protonated form of rotigotine.

In a particular preferred embodiment of the invention the self-adhesive matrix is free of particles that can absorb salts of rotigotine on the TDS/skin interface. Examples of particles that can absorb salts of rotigotine on the TDS/Skin interface include silica. Such particles that can adsorb salts of rotigotine may represent diffusion barriers for the free base form of the drug and may result in the formation of channels inducing some permeability of the self-adhesive matrix for the protonated form of rotigotine. Such embodiments are therefore disadvantageous for practising the invention.

US 8,246,979 B2

5

The self-adhesive matrix of the TDS of the present invention comprises a solid or semi-solid semi-permeable polymer. Usually this polymer will be a pressure sensitive adhesive (PSA) or a mixture of such adhesives. The pressure sensitive adhesive(s) form a matrix in which the active ingredient and the other components of the TDS are incorporated.

The adhesive used in the present invention should preferably be pharmaceutically acceptable in a sense that it is bio-compatible, non-sensitising and non-irritating to the skin. Particularly advantageous adhesives for use in the present invention should further meet the following requirements:

1. Retained adhesive and co-adhesive properties in the presence of moisture or perspiration, under normal temperature variations,
2. Good compatibility with rotigotine, as well as with the further excipients used in the formulation.

Although different types of pressure sensitive adhesive may be used in the present invention, it is preferred to use lipophilic adhesives having both a low drug and low water absorption capacity. Particularly preferably, the adhesives have solubility parameters which are lower than those of rotigotine. Such preferred pressure sensitive adhesives for use in the TDS of the present invention are silicone type pressure sensitive adhesives. Especially preferred pressure sensitive adhesives for use in the TDS of the invention are of the type forming a soluble polycondensed polydimethylsiloxane (PDMS)/resin network, wherein the hydroxy groups are capped with e.g. trimethylsilyl (TMS) groups. Preferred adhesives of this kind are the BIO-PSA silicone pressure sensitive adhesives manufactured by Dow Corning, particularly the Q7-4201 and Q7-4301 qualities. However, other silicone adhesives may also be used.

In a further and especially preferred aspect, two or more silicone adhesives are used as the main adhesive components. It can be advantageous if such a mixture of silicone adhesives comprises a blend of high tack silicone pressure sensitive adhesive comprising polysiloxane with a resin and a medium tack silicone type pressure sensitive adhesive comprising polysiloxane with a resin.

Tack has been defined as the property that enables an adhesive to form a bond with the surface of another material upon brief contact under light pressure (see e.g. "Pressure Sensitive Tack of Adhesives Using an Inverted Probe Machine", ASTM D2979-71 (1982); H. F. Hammond in D. Satas "Handbook of Pressure Sensitive Adhesive Technology" (1989), 2nd ed., Chapter 4, Van Nostrand Reinhold, New York, page 38).

Medium tack of a silicon pressure sensitive adhesive indicates that the immediate bond to the surface of another material is weaker compared to high tack silicon adhesive. The mean resin/polymer ratio is approx. 60/40 for medium tack adhesives, whereas it is approx. 55/45 for high tack adhesives. It is known to the skilled person that both tape and rheological properties are significantly influenced by the resin/polymer ratio (K. L. Ulman and R. P. Sweet "The Correlation of Tape Properties and Rheology" (1998), Information Brochure, Dow Corning Corp., USA).

Such a blend comprising a high and a medium tack silicone type pressure sensitive adhesive comprising polysiloxane with a resin is advantageous in that it provides for the optimum balance between good adhesion and little cold flux. Excessive cold flux may result in a too soft patch which easily adheres to the package or to patient's garments. Moreover, such a mixture seems to be particularly useful for obtaining higher plasma levels. A mixture of the aforementioned

6

Q7-4201 (medium tack) and Q7-4301 (high tack) proved to be especially useful as a matrix for the TDS according to the present invention.

In a further preferred embodiment, the TDS further includes a crystallization inhibitor. Several surfactants or amphiphilic substances may be used as crystallization inhibitors. They should be pharmaceutically acceptable and approved for use in medicaments. A particularly preferred example of such a crystallization inhibitor is soluble polyvinylpyrrolidone, which is commercially available, e.g. under the trademark Kollidon® (Bayer AG). Other suitable crystallization inhibitors include copolymers of polyvinylpyrrolidone and vinyl acetate, polyethyleneglycol, polypropyleneglycol, glycerol and fatty acid esters of glycerol or copolymers of ethylene and vinyl acetate.

The device of the present invention further comprises a backing layer, which is inert to the components of the matrix. This backing layer is a film being impermeable to the active compounds. Such a film may consist of polyester, polyamide, polyethylene, polypropylene, polyurethane, polyvinyl chloride or a combination of the aforementioned materials. These films may or may not be coated with an aluminium film or with aluminium vapour. The thickness of the backing layer may be between 10 and 100 µm, preferably between 15 and 40 µm.

The TDS of the present invention further comprises a protective foil or sheet, which will be removed immediately prior to use, i.e. immediately before the TDS will be brought into contact with the skin. The protective foil or sheet may consist of polyester, polyethylene or polypropylene which may or may not be coated with aluminium film or aluminium vapour or fluoropolymers. Typically the thickness of such a protective foil or sheet ranges from between 50 and 150 µm. So as to facilitate removal of the protective foil or sheet when wishing to apply the TDS, the protective foil or sheet may comprise separate protective foils or sheets having overlapping edges, similar to the kind used with the majority of conventional plasters.

In a preferred embodiment of the present invention, the TDS has a basal surface area of 5 to 50 cm², particularly of 10 to 30 cm². It goes without saying that a device having a surface area of, say, 20 cm² is pharmacologically equivalent to and may be exchanged by two 10 cm² devices or four 5 cm² devices having the same drug content per cm². Thus, the surface areas as indicated herein should be understood to refer to the total surface of all devices simultaneously administered to a patient.

Providing and applying one or several TDS according to the invention has the pharmacological advantage over oral therapy that the attending physician can titrate the optimum dose for the individual patient relatively quickly and accurately, e.g. by simply increasing the number or size of devices given to the patient. Thus, the optimum individual dosage can often be determined after a time period of only about 3 weeks with low side effects.

A preferred content of rotigotine in the TDS according to the invention is in the range of 0.1 to 2.0 mg/cm². Still more preferred are 0.20 to 1.0 mg/cm². If a 7 day patch is desired, higher drug contents will generally be required.

The device used in the present invention is preferably a patch having a continuous adhesive matrix in at least its center portion containing the drug. However, transdermal equivalents to such patches are likewise comprised by the present invention, e.g. an embodiment where the drug is in an inert but non-adhesive matrix in the center portion of the device and is surrounded by an adhesive portion along the edges.

US 8,246,979 B2

7

The TDS according to the present invention is prepared by a manufacturing process, which comprises preparing a rotigotine loaded adhesive, coating, drying or cooling and lamination to get the bulk product, converting the laminate into patch units via cutting, and packaging.

The invention and the best mode for carrying it out will be explained in more detail in the following non-limiting examples.

Invention Example 1

Very Low Salt Content, Small Microreservoirs

252.6 g Rotigotine free base are dissolved in 587.8 g ethanol 100% w/w and mixed with 222.2 g ethanolic solution containing 25% w/w polyvinylpyrrolidone (Kollidon F 90), 0.11% w/w aqueous sodium bisulfite solution (10% w/w), 0.25% ascorbyl palmitate and 0.62% DL- α -tocopherol. To the homogenous mixture 1692.8 g BIO-PSA Q7 4301 (73% w/w), 1691.6 g BIO-PSA Q7 4201 (73% w/w) and 416.3 g petrol ether are added and all components are stirred for at least 1 hour to get a homogenous dispersion.

For manufacture of the patch matrix, the dispersion is coated onto a suitable release liner (for example Scotchpak 1022) and the solvents are continuously removed in a drying oven at temperatures up to 80° C. to get a drug-containing adhesive matrix of 50 g/m² coating weight. The dried matrix film is laminated with a polyester-type backing foil which is siliconized on the inner side and aluminium vapor coated on the opposite side.

The individual patches are punched out of the complete laminate and are sealed into pouches under a nitrogen flow.

The rotigotine contained in the matrix was quantitatively released after 3 hours in the drug dissolution test according to the Paddle over Disk method as described in the USP using the conditions as described above. This result indicates that the obtained TDS was completely free of rotigotine hydrochloride.

The mean size of the microreservoirs in the TDS was approx. 10 μ m with typical sizes in the range of 5 to 35 μ m. A microscope image of the obtained TDS is shown in FIG. 5.

Comparative Example 1

High Salt Content, Small Microreservoirs

2400 g Rotigotine hydrochloride were added to a solution of 272.8 g NaOH in 3488 g ethanol (96%). The resulting mixture was stirred for approximately 10 minutes. Then 379.2 g of sodium phosphate buffer solution (27.6 g Na₂HPO₄·2H₂O) and 53.2 g NaH₂PO₄·2H₂O in 298.5 g water) was added. Insoluble or precipitated solids were separated from the mixture by filtration. The filter was rinsed with 964 g ethanol (96%) to obtain a particle-free ethanolic solution of rotigotine essentially in the form of the free base.

The rotigotine solution (6150 g) in ethanol (30% w/w) was mixed with 407 g ethanol (96%). The resulting solution was mixed with 1738.8 g of an ethanolic solution containing 25 wt. % polyvinylpyrrolidone (Kollidon 90F), 0.11 wt. % aqueous sodium bisulfite solution (10 wt. %), 0.25 wt. % ascorbyl palmitate, and 0.62 wt. % DL- α -tocopherol until homogeneity. To the mixture 13240 g of an amine resistant high tack silicone adhesive (BIO-PSA Q7-4301 mfd. by Dow Corning) (73 wt. % solution in heptane), 13420 g of an amine resistant medium tack silicone adhesive (BIO-PSA® Q7-4201 mfd. by Dow Corning) (72 wt. % solution in heptane), and 3073 g petrol ether were added, and all components were stirred until a homogenous dispersion was obtained.

8

tane), and 3073 g petrol ether were added, and all components were stirred until a homogenous dispersion was obtained.

The dispersion was coated onto a suitable polyester release liner (SCOTCHPAK® 1022) with a suitable doctor knife and the solvents were continuously removed in a drying oven at temperatures up to 80° C. for about 30 minutes to obtain a drug containing adhesive matrix of 50 g/m² coating weight. The dried matrix film was laminated with a polyester-type backing foil (SCOTCHPAK® 1109). The individual patches were punched out of the complete laminate in the desired sizes (e.g. 10 cm², 20 cm², 30 cm²) and sealed into pouches under the flow of nitrogen.

Only approx. 95% of the rotigotine contained in the matrix were released after 3 hours in the drug dissolution test according to the Paddle over Disk method as described in the USP using the conditions as described above. Thus, the obtained TDS contained approx. 5% (w/w) of protonated rotigotine.

The mean size of the microreservoirs in the TDS was approx. 15 μ m with typical sizes in the range of 10 to 20 μ m.

Comparative Example 2

High Salt Content, Large Microreservoirs

150 g Rotigotine hydrochloride were added to a solution of 17.05 g NaOH in 218 g ethanol (96%). The resulting mixture was stirred for approximately 10 minutes. Then 23.7 g of sodium phosphate buffer solution (8.35 g Na₂HPO₄·2H₂O and 16.07 g NaH₂PO₄·2H₂O in 90.3 g water) was added. Insoluble or precipitated solids were separated from the mixture by filtration. The filter was rinsed with 60.4 g ethanol (96%) to obtain a particle-free ethanolic solution of rotigotine essentially in the form of the free base.

The rotigotine solution (346.4 g) in ethanol (35% w/w) was mixed with 36.2 g ethanol (96%). The resulting solution was mixed with 109 g of an ethanolic solution containing 25 wt % polyvinylpyrrolidone (KOLLIDON® 90F), 0.077 wt % aqueous sodium bisulfite solution (10 wt %), 0.25 wt % ascorbyl palmitate, and 0.63 wt % DL- α -tocopherol until homogenous. To the mixture, 817.2 g of an amine resistant high tack silicone adhesive (BIO-PSA® Q7-4301 mfd. by Dow Corning) (74 wt % solution in heptane), 851.8 g of an amine resistant medium tack silicone adhesive (BIO-PSA® Q7-4201 mfd. by Dow Corning) (71 wt % solution in heptane), and 205.8 g petrol ether (heptane) were added, and all components were stirred until a homogenous dispersion was obtained.

The dispersion was coated onto a suitable polyester release liner (SCOTCHPAK® 1022) with a suitable doctor knife and the solvents were continuously removed in a drying oven at temperatures up to 80° C. for about 30 min to obtain a drug-containing adhesive matrix of 50 g/m² coating weight. The dried matrix film was laminated with a polyester-type backing foil (SCOTCHPAK® 1109). The individual patches were punched out of the complete laminate in the desired sizes (e.g. 10 cm², 20 cm², 30 cm²) and sealed into pouches under the flow of nitrogen.

Owing to the large microreservoirs in the TDS' matrix, it was possible to dissolve the rotigotine salts by direct contact with the dissolution medium. Thus, it was not possible to determine the amount of the protonated form of rotigotine. This indicates that the maximum diameter of the microreservoirs was larger than the thickness of the matrix.

The mean size of the microreservoirs in the TDS was approx. 50 μ m with typical sizes in the range of 20 to 90 μ m. A microscope image of the obtained TDS is shown in FIG. 4.

US 8,246,979 B2

9

As rotigotine was released from rotigotine hydrochloride in a manner similar to Comparative Example 1, one may conclude that the obtained TDS also contained 5% (w/w) of rotigotine in its protonated form.

Comparative Example 3

Acrylate-Type Formulation

A mixture of 50.0 g rotigotine hydrochloride and 28.6 g sodium trisilicate in 95 g methyl ethyl ketone was stirred at room temperature for 48 hours. Subsequently, 17.9 g oleic alcohol, 128.6 g of an acrylic-type adhesive solution (51.4% w/w in ethyl acetate; trade name: Durotak® 387-2287 from NATIONAL STARCH & CHEMICAL), 33.0 g of EUDRAGIT® E100 (from ROEHM PHARMA) (50% w/w solution in ethyl acetate) and 45.0 g ethyl acetate were added, and the mass was homogenised mechanically.

The dispersion was coated onto a suitably siliconised process liner (Hostaphan RN 100), and the solvents were evaporated at 50° C. over 30 minutes, thereby obtaining a matrix weight of 60 g/m². The dry film was laminated with a suitable polyester foil (Hostaphan RN 15). Individual patches having a desired size of (e.g. 20 cm²) were punched out of the resulting laminate and sealed into pouches under the flow of nitrogen.

Example 2

In vivo Drug Absorption Test

In order to monitor the absorption of rotigotine by the human skin the following experiment was carried out. The test was performed with the TDS obtained in Example 1 as well as in Comparative Examples 1 and 2.

The plasma concentration time profile at different test times was determined in pharmacokinetic studies involving (A) 14 healthy male persons (TDS of Comparative Examples 2 and 3) or (B) 30 healthy male persons (TDS of Example 1 and Comparative Example 1), respectively. The studies were conducted following an open single-dose randomised (B) two-way or (A) three-way cross-over design.

Individual concentrations of rotigotine were determined by means of liquid chromatography and mass spectroscopy. The lower limit of quantification (LOQ) was 10 µg/ml.

The drug absorption was calculated from the plasma concentration data according to the Wagner-Nelson method (Malcom Rowland, Thomas N. Tozer (Eds.) "Estimation of Adsorption Kinetics from Plasma Concentration Data" in Clinical Pharmacokinetics, pp 480-483, Williams & Wilkins, 1995), 100%=absorption rate after 48 hours; patch application time was 24 hours.

A comparison of the flux across human skin for the different TDS tested is shown in FIGS. 1, 2 and 3.

In FIG. 1 the rotigotine absorption for the sample obtained in Example 1 containing no salt (○) is compared to the sample obtained in Comparative Example 1 containing approx. 5% (w/w) of rotigotine hydrochloride (●). The comparison in FIG. 1 clearly shows that the drug absorption after patch application depends on the residual salt content in the semi-permeable matrix and is significantly improved by reducing the amount of the protonated form of rotigotine present in the matrix.

FIG. 2 shows the impact of the size distribution of the microreservoirs distributed in the semi-permeable matrix by comparing the sample obtained in Comparative Example 1 having a mean microreservoir size of approx. 15 µm and

10

typical sizes between 10 and 20 µm (●) with the sample obtained in Comparative Example 2 having a mean microreservoir size of approx. 50 µm and typical sizes between 20 and 90 µm (▲). From this comparison it can be deduced that reducing the size of the matrix reservoirs significantly increases the flux across the human skin.

A comparison between the TDS of Example 1 (○) and Comparative Example 2 (▲) is shown in FIG. 3. This comparison clearly indicates that the flux across human skin is significantly enhanced by reducing the salt content and decreasing the size of the microreservoirs.

Example 3

In vitro Diffusion Experiment with Transdermal Drug Delivery Systems

The test was performed with a sandwich of consecutively a supportive separator membrane, skin and the TDS. Active substance that has diffused from the TDS through the skin and/or membrane dissolves in an acceptor liquid that continuously passes directly underneath the membrane; the acceptor liquid was collected in tubes in a fraction collector; and the fractions were analysed for their content of rotigotine. The flux of active substance through skin was calculated by correcting for the influence of the separator membrane.

The diffusion cell described in Tanojo et al (Tanojo et al. "New design of a flow through permeation cell for *in vitro* permeation studies across biological membranes" Journal of Controlled Release (1997), 45, 41-47) was used in order to conduct the experiment.

A flask containing the acceptor liquid and the assembled diffusion cells were placed in a temperature-controlled water-bath (32.0±0.5° C.). The acceptor liquid was continuously pumped from the flask through PTFE tubing by a peristaltic pump, passed through the diffusion cells where the diffusion takes place and was then transported via PTFE tubing to test tubes that were placed in a fraction collector.

The required number of disks was punched out from the TDS by using a circular knife. Human epidermis, excised to a thickness of 200-300 µm from fresh donor skin (storage ≤36 hours at 4° C.) with a dermatome (to be referred to as skin) was spread out on laboratory film in petridishes. Using the circular knife the required number of disks was punched out. A disk of membrane was centred on each cell surface. The skin disks were spread out on the membrane disks on the cell surfaces with the aid of forceps. A disk of the TDS is applied to each cell, and the cells were assembled. The experiment was then conducted in a manner similar to the one described in Tanojo et al above.

Afterwards the tubes containing the collected fraction were weighed, and the contents of each tube were analysed using HPLC.

This experiment was carried out for the TDS of Example 1 as well as Comparative Examples 2 and 3.

FIG. 6 shows the *in vitro* skin permeation profile for the TDS of Example 1 (●) compared to the TDS of Comparative Example 2 (○).

FIG. 7 shows the *in vitro* skin permeation profile for the TDS of Example 1 (●) compared to the acrylate TDS of Comparative Example 3 (○).

It is clear from the data obtained that the flux across human skin may be significantly enhanced by controlling the size of the microreservoirs in the TDS while at the same time providing a semi-permeable matrix, which is highly permeable for the free base of rotigotine while being impermeable for its protonated form.

US 8,246,979 B2

11

The invention claimed is:

1. A transdermal delivery system (TDS) comprising a backing layer and a self-adhesive matrix containing rotigotine, wherein the self-adhesive matrix comprises a solid or semi-solid semi-permeable polymer

- (1) wherein rotigotine in its free base form is incorporated, and wherein said free base has been isolated prior to its incorporation into the matrix,
- (2) which comprises a multitude of microreservoirs within the matrix, said microreservoirs containing rotigotine free base,
- (3) which is permeable to the free base of rotigotine,
- (4) which is substantially impermeable to the protonated form of rotigotine, and
- (5) wherein all the microreservoirs have a maximum diameter that is less than the thickness of the matrix; and wherein the backing layer is inert to the components of the matrix.

2. The TDS of claim 1, wherein the microreservoirs have a mean diameter in the range of 0.5 to 20 μm .

3. The TDS of claim 1, wherein the self-adhesive matrix is free of particles that can absorb salts of rotigotine at the TDS/skin interface.

4. The TDS of claim 1, wherein the self-adhesive matrix comprises a silicone pressure sensitive adhesive.

5. The TDS of claim 1, wherein the self-adhesive matrix comprises two or more silicone pressure sensitive adhesives as the main adhesive components.

6. The TDS of claim 5, wherein the two or more silicone pressure sensitive adhesives comprise a blend of a high tack silicone pressure sensitive adhesive comprising polysiloxane with a resin and a medium tack silicone pressure sensitive adhesive comprising polysiloxane with a resin.

12

7. A method for treatment of a patient suffering from a disease treatable with rotigotine, comprising applying the TDS of claim 1 to the skin of the patient.

8. The TDS of claim 1, wherein the microreservoirs additionally contain at least one crystallization inhibitor comprising soluble polyvinylpyrrolidone, a copolymer of polyvinylpyrrolidone and vinyl acetate, polyethylene glycol, polypropylene glycol, glycerol, a fatty acid ester of glycerol and/or a copolymer of ethylene and vinyl acetate.

9. The TDS of claim 8, wherein the at least one crystallization inhibitor comprises soluble polyvinylpyrrolidone.

10. The TDS of claim 1, comprising within the matrix 10^3 to 10^9 microreservoirs per cm^2 of the surface of the matrix.

11. The TDS of claim 1, comprising within the matrix 10^6 to 10^9 microreservoirs per cm^2 of the surface of the matrix.

12. The TDS of claim 1, wherein the microreservoirs have a maximum diameter not greater than 35 μm .

13. The TDS of claim 1, wherein the microreservoirs have a maximum diameter of 2.5 to 30 μm .

14. The TDS of claim 1, wherein any amount of rotigotine salt present comprises less than 5% w/w of the self-adhesive matrix.

15. The TDS of claim 1, wherein any amount of rotigotine salt present comprises less than 2% w/w of the self-adhesive matrix.

16. The TDS of claim 1, wherein any amount of rotigotine salt present comprises less than 1% of the self-adhesive matrix.

17. The TDS of claim 1, wherein the free base of rotigotine has been isolated in solid form prior to the incorporation in the matrix.

18. The TDS of claim 1, wherein the microreservoirs containing rotigotine free base distributed in the matrix have a mean diameter between 1-40% of the thickness of the matrix.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 8,246,979 B2
APPLICATION NO. : 10/623864
DATED : August 21, 2012
INVENTOR(S) : Dietrich Wilhelm Schacht, Mike Hannay and Hans-Michael Wolff

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In the Specifications:

Column 9, line 20, replace “(Hostaphan RN 100)” with --(Hostaphan® RN 100)--.

Column 9, line 23, replace “(Hostaphan RN 15)” with --(Hostaphan® RN 15)--.

Column 10, line 1, replace “(20 Am)” with --(20 μ m)--.

Signed and Sealed this
Third Day of September, 2013



Teresa Stanek Rea
Acting Director of the United States Patent and Trademark Office

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 8,246,979 B2
APPLICATION NO. : 10/623864
DATED : August 21, 2012
INVENTOR(S) : Schacht et al.

Page 1 of 1


It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Title Page:

The first or sole Notice should read --

Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b)
by 1607 days.

Signed and Sealed this
Seventeenth Day of March, 2020



Andrei Iancu
Director of the United States Patent and Trademark Office